

## Antibiotic Resistance in *Salmonella*: A Risk for Tropical Aquaculture

Renata Albuquerque Costa<sup>1</sup>, Fátima Cristiane Teles de Carvalho<sup>2</sup>  
and Regine Helena Silva dos Fernandes Vieira<sup>3</sup>

<sup>1</sup>Doctoral Student in Fisheries Engineering. C.A.P.E.S. Federal University of Ceará

<sup>2</sup>Doctoral Student in Tropical Marine Sciences. Federal University of Ceará

<sup>3</sup>Prof. Dr. Sea Sciences Institute. Federal University of Ceará  
Brazil

### 1. Introduction

Salmonelas are rod-shaped, non-spore-forming Gram-negative facultative anaerobes measuring 0.7-1.5 by 2-5  $\mu\text{m}$ . With the exception of the serovars Gallinarum and Pullorum, salmonelas are motile organisms. They are classified according to morphology and staining pattern and are divided into serotypes and serovars based on their reaction to somatic (O) and flagellar (H) antigens (Bremer et al., 2003). According to Kumar et al.(2003), the genus *Salmonella* has over 2,000 serovars. Two of these—Saintpaul and Newport—have been isolated from seafood (Ponce et al., 2008).

The prevalence of specific *Salmonella* serovars is related to food type. Thus, the serovars Weltevreden and Rissen are predominant in seafoods, as shown by Kumar et al. (2009) in a study on the distribution and phenotypical characterization of *Salmonella* serovars isolated from samples of fish, crustaceans and mollusks from India.

High incidences of *Salmonella* in seafoods have been reported worldwide (Kumar et al., 2010; Asai et al., 2008) in association with outbreaks of fever, nausea, vomiting and diarrhea (Ling et al., 2002). Since *Salmonella* inhabits the intestinal tract of warm-blooded animals, its presence in aquaculture livestock is most likely due to the introduction of fecal bacteria into culture ponds (Koonse et al., 2005). In fact, in a study on *Salmonella* in shrimp, Shabarinath et al. (2007) concluded this pathogen is generally found in rivers and marine/estuarine sediments exposed to fecal contamination.

The quality of aquaculture products may be compromised by exposure to pathogens and biological or chemical contaminants. The latter include chemical agents commonly used in aquaculture, such as veterinary antibiotics, antiseptics and anesthetics. Few antibiotics have been adapted to or developed specifically for use in aquatic organisms. Thus, in Europe several classes of antibiotics may be used in aquaculture, including sulfonamides, quinolones, macrolides, tetracyclines and emamectin. This, however, poses a considerable risk of release of antimicrobials into the environment and eventually of the development of resistance in pathogenic bacteria (Fauconneau, 2002).

The second half of the 20th century saw two major events in the epidemiology of salmonellosis: the appearance of human infections caused by food-borne *S. enteritidis* and by *Salmonella* strains with multiple resistance (Velge et al., 2005). In fact, Angulo et al. (2000) suggested that the factors determining resistance to multiple antibiotics in strains of *S. Typhimurium* DT104 may first have developed in bacteria in the aquaculture environment, possibly as the result of the regular use of antibiotics in fodder.

The present study is a review of the literature on resistant *Salmonella* strains in aquaculture and an assessment of the risk this represents for human health. In addition, information was collected on the incidence of resistant *Salmonella* strains isolated from shrimp farm environments in Northeastern Brazil.

## **2. Methods of isolation, identification and evaluation of antibacterial susceptibility in *Salmonella***

### **2.1 Isolation and identification of *Salmonella***

*Salmonella* may be detected in samples from aquaculture environments using the traditional method described by Andrews and Hammack (2011). The method includes pre-enrichment of 25-g aliquots in lactose broth, selective enrichment in broth (eg, tetrathionate and Rappaport-Vassiliadis or tetrathionate and selenite cystine) and selective plating on MacConkey and Hektoen enteric agar. Typical *Salmonella* colonies grown during the selective enrichment stage are screened biochemically with triple sugar iron agar (TSI), lysine iron agar (LIA) or sulfide indole motility agar (SIM). Serotyping is done with commercially available antisera (Koonse et al., 2005), O:H polyvalent antiserum (Carvalho et al., 2009) or somatic (O), flagellar (H) and capsular (Vi) antisera (Kumar et al., 2009).

In addition, molecular biology techniques may be used for rapid detection of *Salmonella* in foods: TaqMan PCR (Kimura et al., 1999), PCR amplification of a 152-bp segment of the gene *hns* (Kumar et al., 2003), real-time PCR (Malorny et al., 2004), PCR, dot blot hybridization, RAPD and ERIC-PCR (Shabarinath et al., 2007), PCR amplification of the gene *invA* (Upadhyay et al., 2010) and uniplex and multiplex PCR (Raj et al., 2011).

### **2.2 Antibiogram, MIC and plasmid curing**

The phenotypical susceptibility of *Salmonella* to antibiotics may be determined by the method of disk diffusion on Muller-Hinton agar (Kha et al., 2006). When testing salmonellas from aquaculture environments, the selection of antibiotics depends on the origin of the isolates, but usually covers a range of families, including the tetracyclines, sulfonamides, quinolones, macrolides and aminoglycosides (Ponce et al., 2008; Carvalho et al., 2009). The classification of bacteria according to susceptibility or resistance to antibiotics is based on the criteria of the Clinical and Laboratory Standards Institute (CLSI, 2009). The antibacterial resistance index (ARI) may be calculated following the recommendations of Jones et al. (1986). Multiple antibacterial resistance (MAR) may be calculated using the methodology described in Krumpferman (1983).

Antibacterial susceptibility may also be estimated by determining the minimum inhibitory concentration (MIC) based on macrodilution of Mueller-Hinton broth (CLSI, 2009).

*Salmonella* strains with phenotypical profile of antibacterial resistance may be submitted to plasmid curing in Luria-Bertani broth supplemented with acridine orange dye at 100 µg mL<sup>-1</sup>. The method makes it possible to determine whether resistance stems from chromosomal or plasmidial elements (Molina-Aja et al., 2002).

### 2.3 Determination of resistance genes and plasmid profile

Polymerase chain reaction (PCR) has been used to detect genes encoding resistance to tetracycline in *Salmonella* strains from fish farms. Restriction enzymes used in PCR include *Sma*I (for detecting the gene *tetA*), *Sal*I (for *tetC*), *Sph*I (for *tetB*, *tetD* and *tetY*), *Eco*RI (for *G*) and *Nde*II (for *tetE*, *tetH* and *tetI*) (Furushita et al., 2003).

The extraction of plasmidial DNA from salmonelas is usually done by alkaline lysis, as proposed by Birnboim and Doly (1979), with or without modification, or with acidic phenol, as described by Wang and Rossman (1994). For small plasmids, the extraction product may be submitted to electrophoresis in 1% agarose gel following the protocol of Akiyama et al. (2011). The protocol for electrophoresis of mega-plasmid DNA molecules in 1% agarose gel is described in Ponce et al. (2008).

## 3. Results

### 3.1 *Salmonella* in tropical aquaculture

Salmonelas are recognized worldwide as one of the main etiological agents of gastroenteritis in humans. Despite variations in the regulation of microbiological quality of foods around the world, the largest importers of seafoods only buy products completely free from *Salmonella*, based on the claim that salmonelas are not part of the indigenous microbiota of aquatic environments and that, therefore, the presence of salmonelas in aquatic organisms is associated with poor sanitation and inadequate hygiene practices (Dalsgaard, 1998).

Several studies published in the 1990s reported *Salmonella* in shrimp farming environments in tropical countries. Reilly and Twiddy (1992) found *Salmonella* in 16% of their shrimp samples and 22.1% of their pond water and sediment samples collected on farms in Southeast Asia. Weltevreden was the most abundant *Salmonella* serovar identified, followed by Anatum, Wandsworth and Potsdam. According to the authors, the incidence of *Salmonella* was higher in ponds located near urban areas and, not surprisingly, the bacterial load increased during the rainy season. Bhaskar et al. (1995) detected *Salmonella* in 37.5%, 28.6% and 37.4% of shrimp, sediment and water samples, respectively, collected from semi-intensive grow-out ponds in India.

In contrast, despite detecting high indices of thermotolerant and total coliforms, Dalsgaard et al. (1995) found no *Salmonella* in water, sediment and shrimp samples from sixteen different penaeid shrimp farms in Thailand.

Hatha and Rao (1998) reported only one *Salmonella*-positive sample out of 1,264 raw shrimp. They believed the presence of the bacteria was due to pond contamination from different sources, including the use of untreated fertilizer of animal origin. Likewise, Hatha et al. (2003) found the incidence of *Salmonella* to be low in shrimp farm products exported by India.

Koonse et al. (2005) investigated the prevalence of *Salmonella* in six major shrimp-producing countries in Southeast Asia (n=2), Central Asia (n=1), Central America (n=1), North America

(n=1) and the Pacific (n=1). In four of these countries, *Salmonella* was detected in 1.6% of shrimp samples, and two serovars were identified (Paratyphi B var. Java and Weltevreden Z6). The authors highlighted the need to control or eliminate potential sources of fecal matter polluting the water bodies adjacent to the grow-out ponds.

In Brazil, the microbiological quality of shrimp (*Litopenaeus vannamei*) farmed in Ceará was evaluated by Parente et al. (2011) and Carvalho et al. (2009), both of whom detected *Salmonella* in shrimp and water samples (Table 1). The authors associated the presence of salmonelas with discharge of fecal matter into the respective estuaries where the farms are located. The detection of *Salmonella* in estuaries in Ceará is not an isolated finding. Farias et al. (2010) found salmonelas in samples of the bivalve *Tagelus plebeius* collected in the estuary of the Ceará river and identified the serovars Bredeney, London and Muechen. Similar findings were reported by Silva et al. (2003) in a study on *Salmonella* in the oyster *Crassostrea rhizophorae* obtained from natural oyster grounds in the estuary of the Cocó river, on the outskirts of Fortaleza, Ceará.

Country	Sample	N°	Sorovars	Source
Brazil	Water and Shrimp	3	S. ser. Saintpaul e S. ser. Newport	Parente et al. (2011)
Brazil	Fish	30	S. ser. Agona, S. ser. Albany, S. ser. Anatum, S. ser. Brandenburg, S. ser. Bredeney, S. ser. Cerro, S. ser. Enteretidis, S. ser. Havana, S. ser. Infantis, S. ser. Livingstone, S. ser. London, S. ser. Mbandaka, S. ser. Muenchen, S. ser. Newport, S. ser. Saintpaul, S. ser. Thompson, S. ser. O4,5:i:-, S. ser. O4,5:-:1,7, S. O:17	Ribeiro et al., 2010
Brazil	Water, Sediment and Shrimp	23	S. ser. Anatum, S. ser. Newport, S. ser. Soahanina e S. ser. Albany	Carvalho et al. (2009)
Vietnam	Shrimp	29	S. ser. Bovismorbificans, S. ser. Derby, S. ser. Dessau, S. ser. Lexington, S. ser. Schleissheim, S. ser. Tennessee, S. ser. Thompson, S. ser. Virchow, S. ser. Weltevreden, S. ser. II heilbron	Ogasawara et al. (2008)
India	Shrimp	54	S. ser. Bareilly, S. ser. Braenderup, S. ser. Brancaster, S. ser. Derby, S. ser. Kottbus, S. ser. Lindenburg, S. ser. Mbandaka, S. ser. Oslo, S. ser. Rissen, S. ser. Takoradi, S. ser. Typhi, S. ser. Typhimurium, S. ser. Weltevreden, <i>Salmonella</i> VI	Kumar et al. (2009)

\*N°: number of positive samples.

Table 1. *Salmonella* in tropical seafood.

Thus, Shabarinath et al. (2007), who also detected *Salmonella* in shrimp, concluded that since salmonelas inhabit the intestinal tract of warm-blooded animals, their presence in rivers and in marine/estuarine sediments exposed to fecal contamination is not surprising.

Tropical fish species may also be infected with salmonelas (Ponce et al., 2008; Heinitz et al., 2000; Ogbondeminu, 1993); in fact, microorganisms of this genus have recently been associated with farmed catfish (McCoy et al., 2011).

### 3.2 Antimicrobial susceptibility profile of *Salmonella*

The use of antibiotics for prophylaxis in aquaculture not only favors the selection of resistant bacteria in the pond environment, thereby changing the natural microbiota of pond water and sediments, but also increases the risk of transferring resistance genes to pathogens infecting humans and terrestrial animals (Cabello, 2006). Thus, Le and Munekage (2005) reported high levels of drug residues (sulfamethoxazole, trimetoprim, norfloxacin and oxolinic acid) in pond water and sediments from tiger prawn farms in Northern and Southern Vietnam due to indiscriminate use of antibiotics.

According to Seyfried et al. (2010), autochthonous communities in aquatic environments may serve as a reservoir for elements of antibacterial resistance. However, the contribution of anthropic activities to the development of such reserves has not been fully clarified.

Holmström et al. (2003) reported the use, often indiscriminate, of large amounts of antibiotics on shrimp farms in Thailand, and concluded that at a regional scale human health and the environmental balance may be influenced by such practices. Adding to the impact, many of the antibiotics used for prophylaxis in shrimp farming are very persistent and toxic.

Heuer et al. (2009) presented a list of the major antibacterials used in aquaculture and their respective routes of administration: amoxicillin (oral), ampicillin (oral), chloramphenicol (oral, bath, injection), florfenicol (oral), erythromycin (oral, bath, injection), streptomycin (bath), neomycin (bath), furazolidone (oral, bath), nitrofurantoin (oral), oxolinic acid (oral), enrofloxacin (oral, bath), flumequine (oral), oxytetracycline (oral, bath, injection), chlortetracycline (oral, bath, injection), tetracycline (oral, bath, injection) and sulfonamides (oral).

Current aquaculture practices can potentially impact human health in variable, far-reaching and geographically specific ways. On the other hand, the increasing flow of aquaculture products traded on the global market exposes consumers to contaminants, some of which from production areas (Sapkota et al., 2008).

Antibacterial susceptibility in microorganisms associated with aquaculture livestock is an increasingly frequent topic in the specialized literature (Molina-Aja et al., 2002; Peirano et al., 2006; Akinbowale et al., 2006; Costa et al., 2008; Newaja-Fyzul et al., 2008; Dang et al., 2009; Del Cerro et al., 2010; Fernández-Alarcón et al., 2010; Patra et al., 2010; Vieira et al., 2010; Tamminem et al., 2011; Laganà et al., 2011; Millanao et al., 2011; Rebouças et al., 2011; Dang et al., 2011).

In this respect, salmonelas are one of the most extensively investigated groups of intestinal bacteria. Thus, in China salmonelas isolated from fish ponds were resistant to ampicillin

(20%), erythromycin (100%), cotrimoxazole (20%), gentamicin (20%), nalidixic acid (40%), penicillin (100%), streptomycin (20%), sulfanomides (40%), tetracycline (40%) and trimethoprim (20%) (Broughton and Walker, 2009).

Ubeyratne et al. (2008) detected *Salmonella* resistant to erythromycin, amoxicillin and sulfonamides in shrimp (*Penaeus monodon*) farmed in Sri Lanka. Likewise, Ogasawara et al. (2008) found salmonelas resistant to oxytetracycline and chloramphenicol in Vietnamese shrimp samples but concluded ARI values were not as high as in neighboring or developing countries.

Low ARI values were also reported by Boinapally and Jiang (2007) who in a single sample of shrimp imported to the US detected *Salmonella* resistant to ampicillin, ceftriaxone, gentamicin, streptomycin and trimethoprim. This is in accordance with published findings for shrimp in tropical regions, where the major exporters of farmed shrimp are located.

Zhao et al. (2003) evaluated the profile of antibacterial resistance in salmonelas isolated from seafood from different countries and found that most of the resistant bacteria came from Southeast Asia. The authors believe the use of antibiotics in aquaculture, especially in Southeast Asia, favors the selection of resistant *Salmonella* strains which may find their way into the US market of imported foods.

In Brazil, Ribeiro et al. (2010) reported an antibacterial resistance index of 15.1% among salmonelas isolated from an aquaculture system. The *Salmonella* serovars Mbandaka (n=1) and Agona (n=2) were resistant to tetracycline, Albany (n=1) was resistant to sulfamethoxazole-trimethoprim, and London (n=2) was resistant to chloramphenicol. In addition, Carvalho et al. (2009) collected samples from three penaeid shrimp farms in Ceará (Northeastern Brazil) and found *Salmonella* serovars Newport and Anatum to be resistant to tetracycline and nalidixic acid. Water and sediment samples collected in the vicinity of the three farms contained the *Salmonella* serovars Newport, Sohanina, Albany and Anatum, which were likewise resistant to tetracycline and nalidixic acid, suggesting the ponds were contaminated by water drawn from the estuaries.

Bacterial resistance in *Salmonella* may be of either chromosomal or plasmidial nature (Frech e Schwarz, 1999; Mirza et al., 2000; Govender et al., 2009; Tamang et al., 2011; Glenn et al., 2011). In bacteria, the acquisition and diffusion of resistance genes may be influenced by exchanges of DNA mediated by conjugative plasmids and by the integration of resistance genes into specialized genetic elements (Carattoli et al., 2003).

Evidence of plasmidial mediation of antibacterial resistance in *Salmonella* has been available since the 1970s and 1980s (Anderson e Threlfall, 1974; Frost et al., 1982). Thus, Anderson et al. (1977) detected three types of resistance plasmids in *Salmonella* strains from different countries. According to the authors, plasmids of the F<sub>1me</sub> type confer resistance to penicillin, ampicillin and streptomycin, whereas, for example, resistance to furazolidone in all *Salmonella* isolates from Israel was considered to be chromosomal. Mohan et al. (1995) determined the plasmid profile of *Salmonella* strains isolated from different regions in India and found a large diversity of small plasmids (2.7 to 8.3 kb) in strains resistant to ampicillin, chloramphenicol, kanamycin, streptomycin, sulphamethoxazole, tetracycline and trimethoprim.

In one study, salmonelas isolated from food animals were found to carry CMY-2, a plasmid-mediated AmpC-like  $\beta$ -lactamase (Winokur et al., 2001). Doublet et al. (2004) found *florR* (a florfenicol resistance gene) and *bla*<sub>CMY-2</sub> plasmids to be responsible for resistance to wide-spectrum cephalosporines in salmonelas isolated from clinical samples, animals and foods in the US. The authors added that the use of phenicols in animal farming environments may place a selective pressure on organisms and favor the dissemination of *bla*<sub>CMY-2</sub> plasmids. In addition, *florR* is known to confer cross-resistance to chloramphenicol.

Kumar et al. (2010) found evidence that tropical seafood can serve as vehicle for resistant salmonela strains, some of which resistant to as many as four antibiotics (sulfamethizole, carbenicillin, oxytetracycline and nalidixic acid). The authors also identified low-molecular-weight plasmids in the *Salmonella* serovars Braenderup, Lindenburg and Mbandaka.

Six isolates of *Salmonella* serovar Saintpaul from samples of shrimp and fish from India, Vietnam and Saudi Arabia presented one or more resistance plasmids of varying size (2.9 to 86 kb). One of these carried a *Incl1* plasmid (Akiyama et al., 2011).

As discussed above, the indiscriminate use of antibiotics in aquaculture is one of the major causes of the emergence of resistant bacteria in the environment. Several of the mechanisms of resistance in *Salmonella* have been investigated, especially with regard to beta-lactams (Alcaine et al., 2007) and quinolones (Piddock et al., 1998; Piddock, 2002)—two families of antibiotics widely used in aquaculture.

#### 4. Conclusion

The growing incidence of *Salmonella* in tropical aquaculture environments is a worldwide concern which may have local impacts (in the culture area) or global impacts (considering the dynamics of the international seafood market). Human health and environmental balance are further threatened by the emergence of salmonelas resistant to antibiotics employed in farming, in some cases mediated by mobile genetic elements. The elimination of sources of fecal pollution from tropical areas used for aquaculture seems to be the main strategy for minimizing the risk of transference of salmonelas to foods destined for human consumption. As a final consideration, studies should be encouraged on the presence, antibacterial susceptibility and mechanisms of resistance in salmonelas occurring in tropical areas destined for culture of fish, crustaceans and mollusks.

#### 5. References

- Akinbowale OL, Peng H, Barton MD. Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. *Journal of Applied Microbiology*, v. 100, p. 1103-1113, 2006.
- Akiyama T, Khan AA, Cheng CM, Stefanova R. Molecular characterization of *Salmonella* enteric serovar Saintpaul isolated from imported seafood, pepper, environmental and clinical samples. *Food Microbiology*, v. 28, p. 1124-1128, 2011.
- Alcaine SD, Warnick LD, Martin W. Antimicrobial Resistance in Nontyphoidal *Salmonella*. *Journal of Food Protection*, v. 70, n. 3, p. 780-790, 2007.
- Anderson ES, Threlfall EJ, Carr JM, McConnell MM, Smith HR. Clonal distribution of resistance plasmid-carrying *Salmonella typhimurium*, mainly in the Middle East. *Journal of Hygiene*, v. 79, p. 425-448, 1977.


- Anderson ES, Threlfall EJ. The characterization of plasmids in the enterobacteria. *Journal of Hygiene*, v. 72, p. 471-87, 1974.
- Andrews WH, Hammack T. *Salmonella*. In: *Bacteriological Analytical Manual*. U.S. Food and Drug Administration. 2011. Available in: <http://www.fda.gov/Food/ScienceResearch/LaboratoryMethods/BacteriologicalAnalyticalManualBAM/default.htm>.
- Angulo FJ, Griffin PM. Changes in antimicrobial resistance in *Salmonella enterica* serovar Typhimurium. *Emerging Infectious Diseases*, v. 6, n. 4, p. 436-437, 2000.
- Asai Y, Kaneko M, Ohtsuka K, Morita Y, Kaneko S, Noda H, Furukawa I, Takatori K, Harakudo Y. *Salmonella* prevalence in seafood imported into Japan. *Journal of Food Protection*, v. 71, n. 7, p. 1460-1464, 2008.
- Bhaskar N, Setty TMR, Reddy GVS, Manoj YB, Anantha CS, Raghunath BS, Antony JM. Incidence of *Salmonella* in cultured shrimp *Penaeus monodon*. *Aquaculture*, v. 138, p. 257-266, 1995.
- Bhaskar N, Setty TMR, Mondal S, Joseph MA, Raju CV, Raghunath BS, Anantha CS. Prevalence of bacteria of public health significance in the cultured shrimp *Penaeus monodon*. *Food Microbiology*, v. 15, p. 511-519, 1998.
- Birnboim HC, Doly J. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Research*, v. 7, n. 6, p. 1513-1523, 1979.
- Boinapally K, Jiang X. Comparing antibiotic resistance in commensal and pathogenic bacteria isolated from wild-caught South Carolina shrimps vs. farm-raised imported shrimps. *Canadian Journal of Microbiology*, v. 53, n. 7, p. 919-924, 2007.
- Bremer PJ, Fletcher GC, Osborne C. *Salmonella* in seafood. New Zealand Institute for Crop & Food Research Limited, 2003.
- Broughton EL, Walker DG. Prevalence of antibiotic-resistant *Salmonella* in fish in Guangdong, China. *Foodborne pathogens and disease*, v. 6, n. 4, p. 519-521, 2009.
- Cabello FC. Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environmental Microbiology*, v. 8, p. 1137-1144, 2006.
- Carattoli A. Plasmid-mediated antimicrobial resistance in *Salmonella enterica*. *Current Issues in Molecular Biology*, v. 5, p. 113-122, 2003.
- Carvalho FCT, Barreto NSE, Reis CMF, Hofer E, Vieira RHSF. Susceptibilidade antimicrobiana de *Salmonella* spp. Isoladas de fazendas de carcinoculturas no Estado do Ceará. *Revista Ciência Agronômica*, v. 40, n. 4, p. 549-556, 2009.
- CLSI. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; Twentieth Informational Supplement: Supplement M100-S19, Wayne, PA, USA, 2009.
- Costa RA, Vieira GHF, Silva GC, Vieira RHSF, Sampaio SS. Susceptibilidade "in vitro" a antimicrobianos de estirpes de *Vibrio* spp isoladas de camarões (*Litopenaeus vannamei*) e de água de criação destes animais provenientes de uma fazenda de camarões no Ceará - Nota prévia. *Brazilian Journal of Veterinary Research and Animal Science*, v. 45, n. 6, p. 458-462, 2008.
- Dang H, Zhao J, Song L, Chen M, Chang Y. Molecular characterizations of chloramphenicol- and oxytetracycline-resistant bacteria and resistance genes in mariculture waters of China. *Marine Pollution Bulletin*, v. 58, n. 7, p. 987-994, 2009.
- Dang ST, Petersen A, Van Truong D, Chu HT, Dalsgaard A. Impact of medicated feed on the development of antimicrobial resistance in bacteria at integrated pig-fish farms in Vietnam. *Applied and Environmental Microbiology*, v. 77, n. 13, p. 4494-4498, 2011.
- Dalsgaard A. The occurrence of human pathogenic *Vibrio* spp. and *Salmonella* in aquaculture. *International Journal of Food Science and Technology*, v. 33, p. 127-138, 1998.

- Dalsgaard A, Huss HH, H-Kittikun A, Larsen JL. Prevalence of *Vibrio cholerae* and *Salmonella* in a major shrimp production area in Thailand. *International Journal of Food Protection*, v. 28, p. 101-113, 1995.
- Del Cerro A, Márquez I, Prieto JM. Genetic diversity and antimicrobial resistance of *Flavobacterium psychrophilum* isolated from cultured rainbow trout, *Onchorynchus mykiss* (Walbaum), in Spain. *Journal of Fish Diseases*, v. 33, n. 4, p. 285-291, 2010.
- Doublet B, Carattoli A, Whichard JM, White DG, Baucheron S, Chaslus Dancla E, Cloeckeaert A. Plasmid-mediated florfenicol and ceftriaxone resistance encoded by the floR and bla<sub>CMY-2</sub> genes in *Salmonella enterica* serovars Typhimurium and Newport isolated in the United States. *FEMS Microbiology Letters*, v. 233, n. 2, p. 301-305, 2004.
- Fauconneau B. Health value and safety quality of aquaculture products. *Revue de Médecine Vétérinaire*, v. 153, n. 5, p. 331-336, 2002.
- Farias MF, Rocha-Barreira CA, Carvalho FCT, Silva CM, Reis EMF, Costa RA, Vieira RHSF. Condições microbiológicas de *Tagelus plebeius* (Lightfoot 1786) (Mollusca: Bivalvia: Solecurtidae) e da água no estuário do rio Ceará, em Fortaleza-CE. *Boletim do Instituto de Pesca*, v. 36, n. 2, p. 135-142, 2010.
- Fernández-Alarcón C, Miranda CD, Singer RS, López Y, Rojas R, Bello H, Domínguez M, González-Rocha G. Detection of the floR gene in a diversity of florfenicol resistant Gram-negative bacilli from freshwater salmon farms in Chile. *Zoonoses and Public Health*, v. 57, n. 3, p.181-188, 2010.
- Frech G, Schwarz S. Plasmid-encoded tetracycline resistance in *Salmonella enteric* subsp. *enterica* serovars *choleraesuis* and *typhimurium*: identification of complete and truncated Tn1721 elements. *FEMS Microbiology Letters*, v. 176, p. 97-103, 1999.
- Frost JA, Rowe B, Ward LR, Threlfall EJ. Characterization of resistance plasmids and carried phages in an epidemic clone of multi-resistant *Salmonella typhimurium* in India. *Journal of Hygiene*, v. 88, p. 193-204, 1982.
- Furushita M, Shiba T, Maeda T, Yahata M, Kaneoka A, Takahashi Y, Torii K, Hasegawa T, Ohta M. Similarity of tetracycline resistance genes isolated from fish farm bacteria to those from clinical isolates. *Applied and Environmental Microbiology*, v. 69, n. 9, p. 5336-5342, 2003.
- Glenn LM, Lindsey RL, Frank JF, Meinersmann RJ, Englen MD, Fedorka-Cray PJ, Frye JG. Analysis of antimicrobial resistance genes detected in multidrug-resistant *Salmonella enterica* serovar Typhimurium isolated from food animals. *Microbial Drug Resistance*, 2011. DOI:10.1089/mdr.2010.0189.
- Govender N, Smith AM, Karstaedt AS, Keddy KH. Plasmid-mediated quinolone resistance in *Salmonella* from South Africa. *Journal of Medical Microbiology*, v. 58, p. 1393-1394, 2009.
- Hatha AAM, Rao NPB. Bacteriological quality of individually quick-frozen (IQF) raw and cooked ready-to-eat shrimp produced from farm raised black tiger shrimp (*Penaeus monodon*). *Food Microbiology*, v. 15, p. 177-183, 1998.
- Hatha AAM, Maqbool TK, Kumar SS. Microbial quality of shrimp products of export trade produced from aquacultured shrimp. *International Journal of Food Microbiology*, v. 82, p. 213-221, 2003.
- Heinitz ML, Ruble RD, Wagner DE, Tatini SR. Incidence of *Salmonella* in fish and seafood. *Journal of Food Protection*, v. 63, n. 5, p. 579-592, 2000.
- Heuer OE, Kruse H, Grave K, Collignon P, Karunasagar I, Angulo FJ. Human health consequences of use of antimicrobial agents in aquaculture. *Clinical Infectious Diseases*, v. 49, p. 1248-1253, 2009.

- Holmström K, Gräslund S, Wahlström A, Pongshompoo S, Bengtsson BE, Kautsky N. Antibiotic use in shrimp farming and implications for environmental impacts and human health. *International Journal of Food Science and Technology*, v. 38, p. 255-266, 2003.
- Jones JG, Gardener S, Simon BM, Pickup RW. Factors affecting the measurement of antibiotic resistance in bacteria isolated from lake water. *Journal of Applied Microbiology*, v. 60, n. 5, p. 455-462, 1986.
- Khan AA, Cheng CM, Khanh TV, Summage-West C, Nawaz MS, Khan SA. Characterization of class 1 integron resistance gene cassettes in *Salmonella* enteric serovars Oslo and Bareilly from imported seafood. *Journal of Antimicrobial and Chemotherapy*, v. 58, p. 1308-1310, 2006.
- Kimura B, Kawasaki S, Fujii T, Kusunoki J, Ithoh T, Flood SJ. Evaluation of TaqMan PCR assay for detecting *Salmonella* in raw meat and shrimp. *Journal of food protection*, v. 62, n. 4, p. 329-335, 1999.
- Koonse B, Burkhardt III W, Chirtel S, Hoskin GP. *Salmonella* and the sanitary quality of aquacultured shrimp. *Journal of Food Protection*, v. 68, n. 12, p. 2527-2532, 2005.
- Krumperman PH. Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of fecal contamination of foods. *Applied and Environmental Microbiology*, v. 46, p. 165-170, 1983.
- Kumar HS, Sunil R, Venugopal MN, Karunasagar I, Karunasagar I. Detection of *Salmonella* spp. in tropical seafood by polymerase chain reaction. *International Journal of Food Microbiology*, v. 88, p. 91-95, 2003.
- Kumar R, Surendran PK, Thampuran N. Analysis of antimicrobial resistance and plasmid profiles in *Salmonella* serovars associated with tropical seafood of India. *Foodborne Pathogens and Disease*, v. 6, n. 5, p. 621-625, 2009.
- Kumar R, Surendran PK, Thampuran N. Distribution and genotypic characterization of *Salmonella* serovars isolated from tropical seafood of Cochin, India. *Journal of Applied Microbiology*, v. 106, p. 515-524, 2009.
- Kumar R, Surendran PK, Thampuran N. Rapid quantification of *Salmonella* in seafood using real-time PCR assay. *Journal of Microbiology and Biotechnology*, v. 20, n. 3, p. 569-573, 2010.
- Le TX, Munekage Y, Kato S. Antibiotic resistance in bacteria from shrimp farming in mangrove areas. *The Science of the Total Environment*, v. 349, p. 96-105, 2005.
- Laganà P, Caruso G, Minutoli E, Zaccone R, Santi D. Susceptibility to antibiotics of *Vibrio* spp. and *Photobacterium damsela* ssp. *piscicida* strains isolated from Italian aquaculture farms. *The New Microbiologica*, v. 34, n. 1, p. 53-63, 2011.
- Ling ML, Goh KT, Wang GCY, Neo KS, Chua T. An outbreak of multidrug-resistant *Salmonella enterica* subsp. *enterica* serotype Typhimurium, DT104L linked to dried anchovy in Singapore. *Epidemiology and Infection*, v. 128, p. 1-5, 2002.
- Malorny B, Paccassoni E, Fach P, Bunge C, Martin A, Helmuth R. Diagnostic Real-Time PCR for Detection of *Salmonella* in Food. *Applied and Environmental Microbiology*, v. 70, n. 12, p. 7046-7052, 2004.
- McCoy E, Morrison J, Cook V, Johnston J, Eblen D, Guo C. Foodborne agents associated with the consumption of aquaculture catfish. *Journal of Food Protection*, v. 74, n. 3, p. 500-516, 2011.
- Millanao AB, Barrientos MH, Gómez GC, Tomova A, Buschmann A, Dölz H, Cabello FC. Uso inadecuado y excesivo de antibióticos: Salud pública y salmonicultura em Chile. *Revista Médica de Chile*, v. 139, p. 107-118, 2011.

- Mirza S, Kariuki S, Mamun KZ, Beeching NJ, Hart CA. Analysis of plasmid and chromosomal DNA of multidrug-resistant *Salmonella enterica* Serovar Typhi from Asia. *Journal of Clinical Microbiology*, v. 38, n. 4, p. 1449-1452 2000.
- Molina-Aja A, García-Gasca A, Abreu-Grobois A, Bolán-Mejía C, Roque A, Gomez-Gil B. Plasmid profiling and antibiotic resistance of *Vibrio* strains isolated from cultured penaeid shrimp. *FEMS Microbiology Letters*, v. 213, p. 7-12, 2002.
- Mohan VP, Sharma KB, Agarwal DS, Purnima G, Pillai PR. Plasmid profile and phage type of *Salmonella typhimurium* strains encountered in different regions of India. *Comparative Immunology, Microbiology and Infectious Diseases*, v. 18, n. 4, p. 283-290, 1995.
- Newaj-Fyzul A, Mutani A, Ramsubhag A, Adesiyun A. Prevalence of bacterial pathogens and their anti-microbial resistance in Tilapia and their pond water in Trinidad. *Zoonoses and Public Health*, v. 55, n. 4, p. 206-213.
- Ogasawara N, Tran TP, Ly TLK, Nguyen TT, Iwata T, Okatani AT, Watanabi M, Taniguchi T, Hirota Y, Hayashidani H. Antimicrobial susceptibilities of *Salmonella* from domestic animals, food and human in the Mekong delta, Vietnam. *The Journal of veterinary medical science*, v. 70, n. 11, p. 1159-1164, 2008.
- Ogbondeminu FS. The occurrence and distribution of enteric bacteria in fish and water of tropical aquaculture ponds in Nigeria. *Journal of Aquaculture in the Tropics*, v. 8, n. 1, p. 61-66. 1993.
- Parente LS, Costa RA, Vieira GHF, Reis EMF, Hofer E, Fonteles AA, Vieira RHSF. Bactérias entéricas presentes em amostras de água e camarão marinho *Litopenaeus vannamei* oriundos de fazendas de cultivo no Estado do Ceará, Brasil. *Brazilian Journal of Veterinary Research and Animal Science*, v. 48, n. 1, p. 46-53, 2011.
- Patra S, Das TK, Gosh SCh, Sarkar D, Jana BB. Cadmium tolerance and antibiotic resistance of *Pseudomonas* sp. isolated from water, sludge and fish raised in wastewater-fed tropical ponds. *Indian Journal of Experimental Biology*, v. 48, n. 4, p. 383-393, 2010.
- Peirano G, Agerso Y, Aarestrup FM, Reis EMF, Rodrigues DP. Occurrence of integrons and antimicrobial resistance genes among *Salmonella enterica* from Brazil. *Journal of Antimicrobial Chemotherapy*, v. 58, p. 305-309, 2006.
- Piddock LJV, Ricci V, McLaren I, Griggs DJ. Role of mutations in the *gyrA* and *parC* genes of nalidixic-acid-resistant *Salmonella* serotypes isolated from animals in the United Kingdom. *The Journal of antimicrobial chemotherapy*, v. 41, n. 6, p. 635-642, 1998.
- Piddock LJV. Fluoroquinolone resistance in *Salmonella* serovars isolated from humans and food animals. *FEMS Microbiology Reviews*, v. 26, n. 1, p. 3-16, 2002.
- Ponce E, Khan AA, Cheng C-M, Summage-West C, Cerniglia CE. Prevalence and characterization of *Salmonella enterica* serovar Weltevreden from imported seafood. *Food Microbiology*, v. 25, p. 29-35, 2008.
- Raj KT, Jeyasekaran G, Shakila RJ, Thangarani AJ, Sukumar D. Multiplex polymerase chain reaction assay for the detection of *Salmonella enterica* serovars in shrimp in 4 h. *Journal of Bacteriology Research*, v. 3, n. 3, p. 56-62, 2011.
- Rebouças RH, Sousa OV, Lima AS, Vasconcelos FR, Carvalho PB, Vieira RHSF. Antimicrobial resistance profile of *Vibrio* species isolated from marine shrimp farming environments (*Litopenaeus vannamei*) at Ceará, Brazil. *Environmental Research*, v. 111, p. 21-24, 2011.
- Reilly PJ, Twiddy DR. *Salmonella* and *Vibrio cholerae* in brackishwater cultured tropical prawns. *International Journal of Food Microbiology*, v. 16, n. 4, p. 293-301, 1992.
- Ribeiro RV, Reis EMF, Reis CMF, Freitas-Almeida AC, Rodrigues DP. Incidence and antimicrobial resistance of enteropathogens isolated from an integrated aquaculture system. *Letters in Applied Microbiology*, v. 51, p. 611-618, 2010.

- Sapkota A, Sapkota AR, Kucharski M, Burke J, McKenzie S, Walker P, Lawrence R. Aquaculture practices and potential human health risks: Current knowledge and future priorities, v. 34, p. 1215-1226, 2008.
- Seyfried EE, Newton RJ, Rubert IV KF, Pedersen JA, McMahon KD. Occurrence of tetracycline resistance genes in aquaculture facilities with varying use of oxytetracycline. *Microbial Ecology*, v. 59, p. 799-807, 2010.
- Shabarinath S, Sanath Kumar H, Khushiramani R, Karunasagar I, Karunasagar I. Detection and characterization of *Salmonella* associated with tropical seafood. *International Journal of Food Microbiology*, v. 114, n. 2, p. 227-33, 2007.
- Silva A.I.M., Vieira R.H.S.F., Menezes F.G.R., Lima L.N.G.C., Nascimento S.M.M., Carvalho F.C.T. Bactérias fecais em ostras, *Crassostrea rhizophorae*. *Arquivos de Ciências do Mar*, v. 36, p. 63-66, 2003.
- Tamang M.D., Nam H.M., Kim T.S., Jang G.C., Jung S.C., Lim S.K. Emergence of extended-spectrum {beta}-lactamase (CTX-M-15 and CTX-M-14) - producing nontyphoid *Salmonella* with reduced susceptibility to ciprofloxacin among food animals and humans in Korea. *Journal of Clinical Microbiology*, v. 49, n. 7, p. 2671-2675, 2011.
- Tamminen M., Karkman A., Löhmus A., Muziasari W.I., Takasu H., Wada S., Suzuki S., Virta M. Tetracycline resistance genes persist at aquaculture farms in the absence of selection pressure. *Environmental Science & Technology*, v. 45, n. 2, p. 386-391, 2010.
- Upadhyay B.P., Utrarachkij F., Thongshoob J., Mahakunkijcharoen Y., Wongchinda N., Suthienkul O., Khusmith S. Detection of *Salmonella* invA gene in shrimp enrichment culture by polymerase chain reaction. *The Southeast Asian Journal of Tropical Medicine and Public Health*, v. 41, n. 2, p. 426-435, 2010.
- Ubeyratne K.H., Hildebrandt G., Kleer J., Khattiya R., Padungtod P. Microbiological quality of marketed *Penaeus Monodon* shrimps in north western province, Sri Lanka. *Proceedings, The 15<sup>th</sup> Congress of FAVA. OIE Joint Symposium on Emerging Diseases Bangkok, Thailand*, p. P63-P65, 2008.
- Vieira R.H.S.F., Carvalho E.M.R., Carvalho F.C.T, Silva C.M., Sousa O.V., Rodrigues D.P. Antimicrobial susceptibility of *Escherichia coli* isolated from shrimp (*Litopenaeus vannamei*) and pond environment in northeastern Brazil. *Journal of Environmental Science and Health. Part. B, Pesticides, Food Contaminants, and Agricultural Wastes*, v. 45, n. 3, p. 198-203, 2010.
- Velge P., Cloeckaert A., Barrow P. Emergence of *Salmonella enterica* serotype Enteritidis and multiple antibiotic resistance in other major serotypes. *Veterinary Research*, v. 36, p. 267-288, 2005.
- Wang Z., Rossman T.G.. Large-scale supercoiled plasmid preparation by acidic phenol extraction. *Biotechniques*, v. 16, n. 3, p. 460-463, 1994.
- Winokur P.L., Vonstein DL, Hoffman L.J., Uhlenhopp E.K., Doern G.V. Evidence for Transfer of CMY-2 AmpC  $\beta$ -Lactamase Plasmids between *Escherichia coli* and *Salmonella* isolates from food animals and humans. *Antimicrobial Agents and Chemotherapy*, v. 45, n. 10, p. 2716-2722, 2001.
- Zhao S., Datta A.R., Ayers S., Friedman S., Walker R.D., White D.G. Antimicrobial-resistant *Salmonella* serovars isolated from imported foods. *The International Journal of Food Microbiology*, v. 84, p. 87-92, 2003.



## **Salmonella - A Diversified Superbug**

Edited by Mr. Yashwant Kumar

ISBN 978-953-307-781-9

Hard cover, 576 pages

**Publisher** InTech

**Published online** 20, January, 2012

**Published in print edition** January, 2012

Salmonella is an extremely diversified genus, infecting a range of hosts, and comprised of two species: enterica and bongori. This group is made up of 2579 serovars, making it versatile and fascinating for researchers drawing their attention towards different properties of this microorganism. Salmonella related diseases are a major problem in developed and developing countries resulting in economic losses, as well as problems of zoonoses and food borne illness. Moreover, the emergence of an ever increasing problem of antimicrobial resistance in salmonella makes it prudent to unveil different mechanisms involved. This book is the outcome of a collaboration between various researchers from all over the world. The recent advancements in the field of salmonella research are compiled and presented.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Renata Albuquerque Costa, Fátima Cristiane Teles de Carvalho and Regine Helena Silva dos Fernandes Vieira (2012). Antibiotic Resistance in Salmonella: A Risk for Tropical Aquaculture, *Salmonella - A Diversified Superbug*, Mr. Yashwant Kumar (Ed.), ISBN: 978-953-307-781-9, InTech, Available from: <http://www.intechopen.com/books/salmonella-a-diversified-superbug/antibiotic-resistance-in-salmonella-a-risk-for-tropical-aquaculture>

# **INTECH**

open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821